




Contents lists available at ScienceDirect

Clinical Microbiology and Infection

journal homepage: www.clinicalmicrobiologyandinfection.com

Commentary

Human resources estimates and funding for antibiotic stewardship teams are urgently needed

C. Pulcini^{1, 2, 3, *}, C.M. Morel^{4, 5}, E. Tacconelli^{6, 7}, B. Beovic^{3, 8, 9}, K. de With¹⁰,
H. Goossens¹¹, S. Harbarth¹², A. Holmes¹³, P. Howard^{3, 14}, A.M. Morris¹⁵,
D. Nathwani^{16, 17}, M. Sharland¹⁸, J. Schouten^{3, 19}, K. Thursky²⁰, R. Laxminarayan²¹,
M. Mendelson²²

¹ Lorraine University, EA 4360 APEMAC, Nancy, France

² Nancy University Hospital, Infectious Diseases Department, Nancy, France

³ ESCMID Study Group for Antimicrobial stewardship (ESGAP)

⁴ University of Geneva Medical School, Geneva, Switzerland

⁵ London School of Economics, London, United Kingdom

⁶ Infectious Diseases, Internal Medicine 1, DZIF Centre, Tübingen University, Germany

⁷ European Committee on Infection Control (EUCIC)

⁸ University Medical Centre Ljubljana, Slovenia

⁹ Faculty of Medicine, University of Ljubljana, Slovenia

¹⁰ University Hospital Carl Gustav Carus at the TU Dresden, Division of Infectious Diseases, Dresden, Germany

¹¹ Vaccine & Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

¹² Infection Control Program and Division of Infectious Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland

¹³ Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance, Imperial College London, London, UK

¹⁴ Leeds Teaching Hospitals NHS Trust, Leeds, UK

¹⁵ Division of Infectious Diseases, Department of Medicine, Sinai Health System, University Health Network, University of Toronto, Toronto, Canada

¹⁶ Ninewells Hospital and Medical School, Dundee, UK

¹⁷ British Society for Antimicrobial Chemotherapy (BSAC), Birmingham, UK

¹⁸ Paediatric Infectious Diseases Research Group, St George's, University of London, London, UK

¹⁹ IQ Healthcare, Radboud University Medical Centre, Nijmegen, The Netherlands

²⁰ National Centre for Antimicrobial Stewardship, Royal Melbourne Hospital at the Peter Doherty Institute, Melbourne, Australia

²¹ Centre for Disease Dynamics, Economics & Policy, New Delhi, India

²² Division of Infectious Diseases & HIV Medicine, Department of Medicine, Grootte Schuur Hospital, University of Cape Town, Cape Town, South Africa

ARTICLE INFO

Article history:

Received 30 May 2017

Received in revised form

11 July 2017

Accepted 12 July 2017

Available online xxx

Editor: L. Leibovici

Keywords:

Anti-bacterial agents

Antimicrobial

Conservation

Economics

Stewardship

Antibiotic stewardship (AS) teams are essential actors for combating antibiotic-resistant bacteria in healthcare and community settings, and are routinely mentioned in national and international guidelines, recommendations and action plans. Usually, AS teams in resource-rich settings are multidisciplinary, made up of different experts, commonly including infectious diseases (ID) specialists, clinical microbiologists and pharmacists, adequately trained in antibiotic prescribing and stewardship [1]. Some studies conducted in low- and middle-income countries have also shown that community health workers, nurses, village women and others have important roles [1]. Antibiotic stewardship teams are in charge of implementing AS programmes in a specific setting (hospital, community, long-term care facility) or sometimes in different settings, using a cross-sectoral approach [1].

Despite the importance of AS teams in optimizing the management of infections, they remain understaffed or non-existent in most countries [2], and when they do exist, they tend to be focused in hospitals, even though the vast majority of antibiotics are prescribed in the community. A few countries have implemented

* Corresponding author. C. Pulcini, Centre Hospitalier Universitaire de Nancy, Service de Maladies Infectieuses et Tropicales, Hôpitaux de Brabois, allée du Morvan, Vandoeuvre-Lès-Nancy, 54511, France.

E-mail address: celine.pulcini@univ-lorraine.fr (C. Pulcini).

<http://dx.doi.org/10.1016/j.cmi.2017.07.013>

1198-743X/© 2017 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

regulatory measures making hospital stewardship teams mandatory (e.g. Australia, Belgium, Canada, France, Germany, the Netherlands, Norway, UK and the USA), but even these requirements are not always enforced in practice.

Two main actions are required to meet the human resource needs for stewardship across countries; first, there is a need for staffing standards, (e.g. *n* full-time equivalent (FTE) per capita), based on the list of core actions that stewardship teams must implement. Second, a sustainable funding mechanism is needed to ensure that experts in stewardship are employed and have dedicated time for their task. Accepted international norms and standards for infection prevention and control practitioners (per number of hospital beds) have been developed and implemented in some high-income countries, although high heterogeneity still exists at a global level. A recently published literature review concluded that 'an effective infection-control programme in an acute-care hospital must include as a minimum standard at least one full-time specifically trained infection-control nurse per up to 250 beds and a dedicated physician trained in infection control' [3].

To the best of our knowledge, only a handful of countries have established staffing standards for stewardship teams, and these figures only exist for hospitals [4]. The 2016 European Centre for Disease Prevention and Control (ECDC) 'Proposals for European Union (EU) guidelines on the prudent use of antimicrobials in humans' recommended salary support and dedicated time for antimicrobial stewardship hospital-based activities, for example 2–6 FTE per 1000 acute-care beds, based on an expert consensus and citing French and German–Austrian recommendations [5–7]. American colleagues recently suggested 2 FTE ID physicians and 1 FTE ID-trained clinical pharmacist for every 1000 acute care hospital beds in the USA [8]. In 2017, members of ESGAP (the ESCMID Study Group for Antimicrobial stewardship) from 26 different

countries replied to a short e-mail survey about staffing recommendations in their own country; these only existed at national level for hospital-based stewardship teams in Australia, Canada, France, Germany and the Netherlands (Table 1). The observed variation in staffing figures comes from the use of different methodologies to calculate the standards, and also reflects different healthcare systems and organization of care.

Importantly, the list of core activities of AS teams also varies between countries. In some countries, such as the Netherlands, some baseline functions of ID physicians, microbiologists and pharmacists may not be accounted for in the FTE staffing figures, because they are already considered standard of care. Also in Belgium, as of July 2007, all acute-care hospitals and chronic-care hospitals with >150 beds receive financial support from the federal government for hiring a trained antibiotic treatment manager for their AS teams. To this end, an annual budget of € 4.3 million is divided among hospitals according to their number of beds (around € 81 700 per 1000 beds, corresponding to 0.8 FTE per 1000 beds) [9]. The funding, however, only concerns one person supervising implementation of the AS programme, not the whole AS team needed to implement all actions on a daily basis (e.g. ward rounds, systematic advice for specific situations). In contrast, in France, the list of core activities that AS teams have to implement is much longer, and includes supervision of the programme as well as actions such as daily advice to prescribers; hospital-based AS teams are also expected to participate in AS activities in outpatient regional networks [7]. A list of core AS activities needs to be agreed upon globally, in addition to the list of basic resources (e.g. diagnostics, pharmacy services) that must be in place for an AS programme to function properly.

The need for a global funding mechanism to address human resources for stewardship, infection prevention and other non-drug

Table 1
Staffing recommendations available at country level for antibiotic stewardship teams in hospitals

Country	How these standards were defined	Staffing standards	Reference
Australia	Antimicrobial stewardship in Australian hospitals (Second edition due 2017)	There is no consensus on staffing recommendations in Australia within the national accreditation standards. Recommendations from 2011: clinicians in hospitals with existing programmes suggest that for every 100 acute beds, at least 10 hours (0.3 FTE) of senior pharmacist and 3.5 hours (0.1 FTE) of lead clinician time per week should be dedicated to AMS activities. Minimum for the team: 4 FTE per 1000 acute-care beds. For rural regional hospitals access to experts through networks or telehealth is recommended.	[12]
Canada	Based on an environmental scan, survey of the medical literature, and expert opinion of the Antimicrobial Stewardship and Resistance Committee	Core Team Members (minimum recommended): total of 4.9 FTE/1000 acute-care beds Physician: 1.0 FTE per 1000 acute-care beds Pharmacist: 3.0 FTE per 1000 acute-care beds Project/Programme Administrative and Coordination Support: 0.5 FTE per 1000 acute-care beds Data Analyst: 0.4 FTE per 1000 acute-care beds	[13]
Austria and Germany	Guideline by the German Society for Infectious Diseases, based on the literature and expert advice	Antibiotic stewardship team: minimum of 2 FTE per 1000 beds. The team should consist of at least one infectious diseases physician (or clinician with infectious diseases training) and an experienced clinical pharmacist/hospital pharmacist, as well as a specialist in microbiology.	[6]
France	Nationwide survey in 65 hospitals, conducted in 2015 by a Task force on antimicrobial resistance coordinated by the Ministry of Health	Optimal standards for the whole antibiotic stewardship team: —3.6 FTE per 1000 acute-care beds for infection specialists (medical doctors, ideally infectious diseases specialists) —2.5 FTE per 1000 acute-care beds for pharmacists —and 0.6 FTE per 1000 acute-care beds for microbiologists i.e. a total of 6.7 FTE per 1000 acute-care beds for the whole antibiotic stewardship team.	[7]
The Netherlands	National consensus procedure	Standards for the whole antibiotic stewardship team: <i>Start-up phase: Optimal standards</i> Hospital <300 beds: 100 hours one time + 0.87 FTE per year Hospital 300–750 beds: 100 hours one time + 1.2 FTE per year Hospital >750 beds: 100 hours one time + 1.53 FTE per year <i>Consolidation phase: Minimum standards</i> Hospital <300 beds: 1.25 FTE per year Hospital 300–750 beds: 2.14 FTE per year Hospital >750 beds: 3.0 FTE per year	[14]

Abbreviation: FTE, full-time equivalent.

Box 1**Steps urgently needed to move forward**

1. Draw up minimum staffing standards for antibiotic stewardship teams that can be used globally. The simplest place to start is probably hospital-based stewardship activities. The first step would be to come up with a list of core antibiotic stewardship activities that would be applicable worldwide, based on a literature review and consensus procedure involving an international panel of experts. The next step would be to estimate the human resources needed to complete all these activities, based on data coming from existing fully implemented and successful programmes in a representative sample of hospitals, completed by experts' opinion.
2. Apply costs to these standards for a sample of low-, middle- and high-income countries.
3. Identify potential funding sources to support health systems to cover these costs.

development measures has been highlighted [10], but remains elusive. Although more keenly felt in low- and middle-income countries, we believe that it is pertinent to all countries, as sustainable protected time for stewardship activities is often lacking. The role of the funding would be to train and retain core healthcare professionals and provide essential support staff (e.g. administrative staff and data support). Funding for AS should not come from the assumed cost savings deriving from the lower drugs expenses, but rather should be an inherent part of patient safety and healthcare quality-related spending.

This is especially urgent given the increasing difficulty to bring new antibiotics to market and the dramatic sums of public money that will likely be needed to do so. Although the cost of stewardship will seem very small relative to these research and development investments, it must not be an afterthought [11]. Indeed it is essential that future stewardship activity has its own earmarked funding stream within any new antibiotic incentive scheme if we are to have any chance of protecting these new drugs if and when they make it to market.

In conclusion, it is time we focused more on optimizing the use of antibiotics, globally. We need further studies to identify the minimum international staffing standards for stewardship teams. This should go beyond hospitals, and should include also the community setting and long-term care facilities. This call recognizes the fact that as standards are put in place, they must take into account different models of stewardship delivery, where both traditional role players and others such as non-specialist pharmacists, nurses and community health workers can be part of AS teams. We call for the development of global estimates of funding needs for basic and core antibiotic stewardship activities across all healthcare settings (Box 1). This is the first step in a long road towards a global funding mechanism to ensure compliance with the standards.

Transparency declaration

There are no conflicts of interest to declare for any of the authors.

Acknowledgements

We thank all the ESGAP members who have kindly replied to our questionnaire. We also thank the ESGAP Executive Committee members who reviewed this manuscript (Guillaume Béraud and Oliver J. Dyar).

Funding

C. Morel has received support from the Innovative Medicines Initiative Joint Undertaking under grant agreement no. 115618 (Driving re-investment in R&D and responsible antibiotic use—DRIVE-AB—www.drive-ab.eu), resources of which comprise a financial contribution from the European Union's Seventh Framework Programme (FP7/2007-2013) and EFPIA companies' in-kind contribution. There is no funding to declare for any of authors.

References

- [1] Dyar OJ, Tebano G, Pulcini C. Managing responsible antimicrobial use: perspectives across the healthcare system. *Clin Microbiol Infect* 2017;23:441–7.
- [2] Howard P, Pulcini C, Levy Hara G, West RM, Gould IM, Harbarth S, et al. An international cross-sectional survey of antimicrobial stewardship programmes in hospitals. *J Antimicrob Chemother* 2015;70:1245–55.
- [3] Zingg W, Holmes A, Dettenkofer M, Goetting T, Secci F, Clack L, et al. Hospital organisation, management, and structure for prevention of health-care-associated infection: a systematic review and expert consensus. *Lancet Infect Dis* 2015;15:212–24.
- [4] Dickstein Y, Nir-Paz R, Pulcini C, Cookson B, Beović B, Tacconelli E, et al. Staffing for infectious diseases, clinical microbiology and infection control in hospitals in 2015: results of an ESCMID member survey. *Clin Microbiol Infect* 2016;22: 812.e9–812.e17.
- [5] ECDC. Proposals for EU guidelines on the prudent use of antimicrobials in humans. 2017. Available at: http://ecdc.europa.eu/en/publications/_layouts/Publication_DispForm.aspx?List=4f55ad51-4aed-4d32-b960-af70113dbb90&ID=1643.
- [6] de With K, Allerberger F, Amann S, Apfalter P, Brodt HR, Eckmanns T, et al. Strategies to enhance rational use of antibiotics in hospital: a guideline by the German Society for Infectious Diseases. *Infection* 2016;44:395–439.
- [7] Le Coz P, Carlet J, Roblot F, Pulcini C. Human resources needed to perform antimicrobial stewardship teams' activities in French hospitals. *Med Mal Infect* 2016;46:200–6.
- [8] Spellberg B, Bartlett JG, Gilbert DN. How to pitch an antibiotic stewardship program to the hospital C-Suite. *Open Forum Infect Dis* 2016;3: ofw210.
- [9] Van Gastel E, Balligand E, Costers M, Magerman K. Antibiotic management teams in Belgian hospitals: continued improvement in the period from 2007 to 2011. *Eur J Clin Microbiol Infect Dis* 2015;34:673–7.
- [10] Mendelson M, Dar OA, Hoffman SJ, Laxminarayan R, Mpundu MM, Røttingen JA. A global antimicrobial conservation fund for low- and middle-income countries. *Int J Infect Dis* 2016;51:70–2.
- [11] Laxminarayan R. Antibiotic effectiveness: balancing conservation against innovation. *Science* 2014;345:1299–301.
- [12] <https://www.safetyandquality.gov.au/wp-content/uploads/2011/01/Antimicrobial-stewardship-in-Australian-Hospitals-2011.pdf>.
- [13] <https://www.ammi.ca/?ID=126&Language=ENG>.
- [14] <http://www.ateams.nl/documenten/>.